Exercise 12.1

Q.1 Prove that the centre of a circle is on the right bisectors of each of its chords.

Given

A, B, C are the three non-collinear points.

Required: To find the centre of the circle passing through A,B,C


Construction

Join B to C, A take \overrightarrow{PQ} is right bisector of \overline{AB} and \overline{RS} right bisector of BC, they intersect at O.

Join O to A, O to B, O to C.

.: O is the centre of circle.

Proof



↑R

Statements	Reasons
$\overline{OB} \cong \overline{OC}$ (i)	O is the right bisector of \overline{BC}
$\overline{OA} \cong \overline{OB}$ (ii)	O is the right bisector of \overline{AB}
$\overline{OA} = \overline{OB} = \overline{OC}$	From (i) and (ii)
Hence is equidistant from the A,B,C	
$\therefore O$ is center of circle which is required	

Q.2 Where will the center of a circle passing through three non-collinear points? And Why? Given

A.B.C are three non collinear points and circle passing through these points.

To prove

Find the center of the circle passing through vertices A, B and C.

Construction

- (i) Join B to A and C.
- (ii) Take \overrightarrow{QT} right bisector of \overline{BC} and take also \overrightarrow{PR} right bisector of \overline{AB} .

 \overrightarrow{PR} and \overrightarrow{QT} intersect at point O. joint O to A,B and C. O is the center of the circle.

Statements	Reasons
\overline{QO} is right bisector \overline{BC}	
$\overline{OB} \cong \overline{OC}$ (i)	
\overline{PO} is right bisector of \overline{AB}	
$\overline{OA} \cong \overline{OB}$ (ii)	
So	F (') 1 ('')
$\overline{OA} \cong \overline{OC} \cong \overline{OB}$	From (i) and (ii)
:. It is proved that O is the center of the circle.	

Q.3 Three village P,Q and R are not on the same line. The people of these villages want to make a children park at such a place which is equidistant from these three villages. After fixing the place of children park prove that the park is equidistant from the three villages.

Given

P,Q,R are three villages not on the same straight line.

To prove

The point equidistant from P,R,Q.

Construction

- (i) Joint Q to P and R.
- (ii) Take \overrightarrow{AB} right bisector of \overrightarrow{PQ} and \overrightarrow{CD} right bisector of \overrightarrow{QR} . \overrightarrow{AB} and \overrightarrow{CD} intersect at O.
- (iii) Join 0 to P, Q, R

 The place of children part at point O.

Proof

Statements	Reasons
$\overline{OQ} \cong \overline{OR}$ (i)	O is on the right bisector of \overline{QR}
$\overline{OP} \cong \overline{OQ}$ (ii)	O is on the right bisector of \overline{PQ}
$\overline{OP} \cong \overline{OQ} \cong \overline{OR}$ (iii)	From (i) and (ii)
$\therefore O$ is on the bisector of $\angle P$	
Hence \overline{PO} is bisector of $\angle P$	

O is equidistant from P,Q and R

Theorem 12.1.3

The right bisectors of the sides of a triangle are concurrent.

Given


 ΔABC

To prove

The right bisectors of \overline{AB} , \overline{BC} and \overline{CA} are concurrent.

Construction

Draw the right bisectors of \overline{AB} and \overline{BC} which meet each other at the point O. Join O to A, B and C.



Theorem 12.1.4

Any point on the bisector of an angle is equidistant from its arms.

Given

A point P is on \overrightarrow{OM} , the bisector of $\angle AOB$


To Prove

 $\overline{PQ} \cong \overline{PR}$ i.e P is equidistant from \overrightarrow{OA} and \overrightarrow{OB}

Construction

Draw $\overrightarrow{PR} \perp \overrightarrow{OA}$ and $\overrightarrow{PQ} \perp \overrightarrow{OB}$

Proof



Theorem 12.1.5 (Converse of Theorem 12.1.4)

Any point inside an angle, equidistant from its arms, is on the bisector of it.

Given

Any point P lies inside $\angle AOB$, such that $\overline{PO} + \overline{OP} + \overline$


 $\overline{PQ} \cong \overline{PR}$, where $\overline{PQ} \perp \overrightarrow{OB}$ and $\overline{PR} \perp \overrightarrow{OA}$


To prove

Point P is on the bisector of $\angle AOB$

Construction

Join P to O





Exercise 12.2

Q.1 In a quadrilateral ABCD $\overline{AB}\cong \overline{BC}$ and the right bisectors of $\overline{AD},\overline{CD}$ meet each other at point N. Prove that \overline{BN} is a bisector of $\angle ABC$

Given

In the quadrilateral ABCD

$$\overline{AB} \cong \overline{BC}$$

 \overline{NM} is right bisector of \overline{CD}

 \overline{PN} is right bisector of \overline{AD}


They meet at N

To prove

 \overline{BN} is the bisector of angle ABC


Construction join N to A,B,C,D

Proof



11001	
Statements	Reasons
$\overline{ND} \cong \overline{NA}$ (i)	N is an right bisector of \overline{AD}
$\overline{ND} \cong \overline{NC}$ (ii)	N is on right bisector of \overline{DC}
$\overline{NA} = \overline{NC}$ (iii)	from (i) and (ii)
$\Delta BNC \leftrightarrow \Delta ANB$	
$\overline{NC} = \overline{NA}$	Already proved (from iii)
$\overline{AB} \cong \overline{CB}$	Given
$\overline{BN} \cong \overline{BN}$	Common
$\therefore \Delta BNA \cong \Delta BNC$	$S.S.S \cong S.S.S$
Hence $\angle ABN \cong \angle NBC$	Corresponding angles of congruent triangles
Hence \overline{BN} is the bisector of $\angle ABC$	× 1/2

Q.2 The bisectors of $\angle A, \angle B$ and $\angle C$ of a quadrilateral ABCP meet each other at point O. Prove that the bisector of $\angle P$ will also pass through the point O.



Given

ABCP is quadrilateral. \overline{AO} , \overline{BO} , \overline{CO} are bisectors of $\angle A$, $\angle B$ and $\angle C$ meet at point O.

To prove

 \overline{PO} is bisector of $\angle P$

Construction:

Join P to O.

Draw $\overline{OQ} \perp \overline{AP}$, $\overline{ON} \perp \overline{PC}$ and $\overline{OL} \perp \overline{AB}$, $\overline{OM} \perp \overline{BC}$

Proof:

Statements	Reasons
$\overline{OM} \cong \overline{ON}$ (i)	O is on the bisector of $\angle C$
$\overline{OL} \cong \overline{OM}$ (ii)	O is on the bisector of $\angle B$
$\overline{OL} \cong \overline{OQ}$ (iii)	O is on the bisector of $\angle A$
$\overline{OQ} \cong \overline{ON}$	From i, ii, iii
Point O lines on the bisector of $\angle P$	
$\therefore \overline{OP}$ is the bisector of angle P	

Q.3 Prove that the right bisector of congruent sides of an isosceles triangle and its altitude are concurrent.

Given

 ΔABC

 $\overline{AB} \cong \overline{AC}$ due to isosceles triangle \overline{PM} is right bisector of \overline{AB}


 \overline{QN} is right bisector of \overline{AC}

 \overrightarrow{PM} and \overrightarrow{QN} intersect each other at point O

Required

The altitude of $\triangle ABC$ lies at point O

Join A to O and extend it to cut \overline{BC} at D.



 $\angle BAD \cong \angle CAD$ $\Delta BAD \cong \Delta CAD$ $\angle ODB \cong \angle ODC$ $m\angle ODM + m\angle ODC = 180^{\circ}$

 $\therefore \overline{AD} \perp \overline{BC}$

Point 0 lies on altitude \overline{AD}

Proved from (i) $S.A.S \cong S.A.S$

Each angle is 90° (Given)

Supplementary angle

Q.4 Prove that the altitudes of a triangle are concurrent.

Given

In ΔABC

AD, BE, CF are its altitudes


i.e $\overline{AD} \perp \overline{BC}, \overline{BE} \perp \overline{AC}, \overline{CF} \perp \overline{AB}$

Required AD, BE and CF are concurrent



Passing through A, B, C take

 $\overline{RQ} \| \overline{BC}, \overline{RP} \| \overline{AC} \text{ and } \overline{QP} \| \overline{AB} \text{ respectively forming a } \Delta PQR$



Statements	Reasons
$\overline{\mathrm{BC}} \ \overline{\mathrm{AQ}} \ $	Construction
$\overline{AB} \ \overline{QC}$	Construction
∴ ABCQ is a ^{gm}	
Hence $\overline{AQ} = \overline{BC}$	9,
Similarly $\overline{AB} \cong \overline{QC}$	
Hence point A is midpoint RQ	
And $\overline{AD} \perp \overline{BC}$	Given
$ \overline{BC} \overline{RQ}$	Opposite sides of parallelogram ABCQ
$\overline{AD} \ \overline{RQ} \ $	
Thus $\overline{AD} \perp$ is right bisector of \overline{RQ}	
similarly BE is a right bisector of RP and	
CF is right bisector of PQ	
$\therefore \perp^{s} \overline{AD}, \overline{BE}, \overline{CF}$ are right bisector of sides of ΔPQR	
$\therefore \overline{AD}, \overline{BE}$ and \overline{CF} are	
Concurrent	

Theorem12.1.6

The bisectors of the angles of a triangle are concurrent

Given

 ΔABC


To Prove

The bisector of $\angle A$, $\angle B$, and $\angle C$ are concurrent

Construction:

Draw the bisectors of $\angle B$ and $\angle C$ which intersect at point I. From I, draw


 $\overline{\text{IF}} \perp \overline{\text{AB}}, \overline{\text{ID}} \perp \overline{\text{BC}} \text{ and } \overline{\text{IE}} \perp \overline{\text{CA}}$



Statements	Reasons	
$\overline{ID} \cong \overline{IF}$	(Any point on bisector of an angle is equidistance from its arms.	
Similarly		
ID ≅ IE_		
∴ IE≅IF	Each ≅ ID	
So the point I is on the bisector of $\angle A \dots (i)$		
Also the point I is on the bisectors of ZABC and ZBCA (ii)	Construction	
Thus the bisector of $\angle A$, $\angle B$ and $\angle C$ are concurrent at I	{From (i) and (ii)}	
	10)	

Exercise 12.3

Q.1 Prove that the bisectors of the angles of base of an isosceles triangle intersect each other on its altitude.



Given


 ΔABC

 $\overline{AB} = \overline{AC}$ Due to isosceles triangle

Bisect $\angle B$ and $\angle C$ to intersect at point O Join A to D and extend to BC at D \overline{AD} is the altitude of $\triangle ABC$ $\overline{AD} \perp \overline{BC}$

Statements	Reasons
In ΔABC	
$\overline{AB} \cong \overline{AC}$	Given
$\angle B \cong \angle C$	Due to isosceles triangle opposite angle are congruent
$\frac{1}{2} \mathbf{m} \angle \mathbf{B} = \frac{1}{2} \mathbf{m} \angle \mathbf{C}$	Dividing both side by 2
$\angle 1 \cong \angle 3$	
$\Delta ABO \leftrightarrow \Delta ACO$	
$\underline{AO} = \underline{AO}$	
$\overline{AB} = \overline{AC}$	
BO≅CO	Given
$\Delta ABO \cong \Delta ACO$	Due to isosceles triangle
$\Delta ABD \leftrightarrow \Delta ACD$	
$\overline{AD} \cong \overline{AD}$	
∠ 7 ≅ ∠ 8	
$\overline{AB} \cong \overline{AC}$	
$\Delta ABD \cong \Delta ACD$	
∠5+∠6 = 180	
$\angle 5 = \angle 6 = 90^{\circ}$	
So $\overline{AD} \perp \overline{BC}$	Supplementary angles
AD Passes from point O	

Q.2 Prove that the bisectors of two exterior and third interior angle of a triangle are concurrent



Given

 ΔABC

Exterior angles are $\angle ABQ$ and $\angle BAP$ \overrightarrow{AT} and \overrightarrow{BS} intersect each other at point O therefore join O to C

Draw the angle bisecter of C

Construction

 $\overrightarrow{OM} \perp \overrightarrow{CQ}, \overrightarrow{OL} \perp \overrightarrow{AB}, \overrightarrow{ON} \perp \overrightarrow{CP}$

Statements	Reasons
$\overline{ON} \cong \overline{OM}$ (i)	
$\overline{OL} \cong \overline{OM}$ (ii)	
$\overline{ON} \cong \overline{OL}$	
Hence Angle Bisector of C	Comparing equation (i) and (ii)
$i,e \angle 1 \cong \angle 2$	

Review Exercise 12

Q.1 Which of the following are true and which are false?

Bisection means to divide into two equal parts (i)

(True)

- Right bisection of line segment means to draw perpendicular which passes through the (ii) midpoint of line segment (True)
- Any point on the right bisector of a line segment is not equidistant from its end points (iii)

(False)

Any point equidistant from the end points of a line segment is on the right bisector of it (iv)

> (True) (False)


- **(v)** The right bisectors of the sides of a triangle are not concurrent
- The bisectors of the angles of a triangle are concurrent (True) (vi)
- Any point on the bisector of an angle is not equidistant from its arms (vii) (False)
- Any point inside an angle equidistant from its arms, is on the bisector of it (True) (viii)

If \overrightarrow{CD} is right bisector of line segment \overrightarrow{AB} , then **Q.2**

- (i) $m\overline{OA} =$
- (ii) $m\overline{AQ} =$


Solution

- (i) $m\overline{OA} = m\overline{OB}$
- (ii) $m\overline{AQ} = m\overline{BQ}$



Define the following Q.3


Right Bisector of a Line Segment (i)



A line *l* is called a right bisector of a line segment if *l* is perpendicular to the line segment and passes through its midpoint.

(ii) Bisector of an Angle

A ray BP is called the bisector of m \(\times ABC \), if P is a point in the interior of the angle and $m \angle ABP = m \angle PBC$.



The given triangle ABC is equilateral triangle and \overline{AD} is bisector of angle A, then find, **Q.4** the values of unknown x^0 , y^0 and z^0 .

Solution

In equilateral triangle all side are equal to each and there angle of the triangle equal to 60°. So

$$\angle B = z^o = 60^o$$

$$\overline{AD}$$
 is the bisector of $\angle A$

$$\angle A = 60^{\circ}$$

:. When angle A is bisected

$$x^{\circ} = y^{\circ}$$

$$x^{\circ} = \frac{1}{2}m\angle A$$


$$=\frac{1}{2}\times60^{\circ}$$

$$x^{\circ} = 30^{\circ}$$

$$y^{\circ} = 30^{\circ}$$

$$(x^{\circ} = y^{\circ})$$

$$y^{\circ} = 30^{\circ}$$
 (: $x^{\circ} = y^{\circ}$)
So $x^{\circ} = y^{\circ} = 30^{\circ}$



In the given congruent triangle LMO and LNO find the unknowns x and m given **Q.5**

$$\Delta LMO \cong \Delta LNO$$

$$m\overline{\text{LM}} = m\overline{\text{LN}}$$

$$2x + 6 = 18$$

$$2x = 18 - 6$$


$$2x = 12$$

$$x = \frac{12^{6}}{2}$$

$$x = 6$$
 Unit

$$m\overline{\text{MO}} = m\overline{\text{ON}}$$

$$\therefore m = 12 \text{ unit}$$



CD is right bisector of the line segment AB **Q.6**

If $m\overline{AB} = 6cm$ then find the $m\overline{AL}$ and $m\overline{LB}$ (i)


Solution

L is the midpoint of \overline{AB}

$$\therefore m\overline{AL} = m\overline{LB}$$

$$m\overline{AL} = \frac{1}{2}mAB = \frac{1}{2} \times 6$$

So
$$m\overline{AL} = 3$$
cm



$$m\overline{LB} = 3$$
cm $\left(: m\overline{AL} = m\overline{LB} \right)$

(ii) If $m\overline{BD} = 4cm$ then find $m\overline{AD}$

 $m\overline{AD} = m\overline{BD}$ (Any point on the right bisector of a line segment is equidistant from its end points.)

$$m\overline{AD} = 4$$

$$m\overline{AD} = 4cm$$



Unit 12: Line Bisectors and Angle Bisectors

Overview

Right Bisector of a line segment:

Right bisection of a line segment means to draw a perpendicular at the mid-point of line segment.

Bisector of an angle:

Bisection of an angle means to draw a ray to divide the given angle into two equal parts.

Theorem 12.1.1

Statement:

Any point on the right bisector of a line segment is equidistant from its end points.

Given

A line \overrightarrow{LM} intersects the line segment AB at the point C Such that $\overrightarrow{LM} \perp \overline{AB}$ and $\overline{AC} \cong \overline{BC}$. P is a point on \overrightarrow{LM}




$$\overline{PA} \cong \overline{PB}$$

Construction

Join P to the point A and B

Proof



Statements	Reasons
In $\triangle ACP \leftrightarrow \triangle BCP$	
$\overline{AC} \cong \overline{BC}$	Given
$\angle ACP \cong \angle BCP$	Given $\overline{PC} \perp \overline{AB}$, so that each \angle at $C = 90^{\circ}$
$\overline{PC} \cong \overline{PC}$	Common
$\therefore \Delta ACP \cong \Delta BCP$	S.A.S Postulate
Hence $\overline{PA} \cong \overline{PB}$	(Corresponding sides of congruent triangles)


Theorem 12.1.2

{Converse of Theorem 12.1.1}

Any point equidistant from the end points of a line segment is on the right bisector of it.

Given

 \overline{AB} is a line segment. Point P is such that $PA \cong PB$



To prove

The point P is the on the right bisector of \overline{AB}

Construction

Join P to C, the midpoint of \overline{AB}

Statements	Reasons
In $\triangle ACP \leftrightarrow \triangle BCP$	
$\overline{PA} \cong \overline{PB}$	Given
$\overline{PC} \cong \overline{PC}$	Common
$\overline{AC} \cong \overline{BC}$	Construction
$\therefore \Delta ACP \cong \Delta BCP$	$S.S.S \cong S.S.S$
$\angle ACP \cong \angle BCP$ (i)	Corresponding angles of congruent triangles
But $m\angle ACP + m\angle BCP = 180^{\circ}$ (ii)	Supplementary angles
$\therefore m \angle ACP = m \angle BCP = 90^{\circ}$	From (i) and (ii)
i.e $\overline{PC} \perp \overline{AB}$ (iii)	$m\angle ACP = 90^{\circ} (Proved)$
Also $\overline{CA} \cong \overline{CB}$ (iv)	Construction
$\therefore \overline{PC}$ is a right bisector of \overline{AB}	from (iii) and (iv)
i.e. the point P is on the right bisector of \overline{AB}	